

CURRICULUM VITAE

Thi-Sau Migone

Birth: Saigon 12/12/1968

Nationality: Italian

Marital status: married, one child

2001-date Senior Scientist I
Department of Pre-clinical Development,
Immunology Section,
Human Genome Sciences, Inc.
9410 Key West Avenue,
Rockville MD, 20850

1999-2001 Scientist
Department of Pre-clinical Development,
Immunology Section,
Human Genome Sciences, Inc.
9410 Key West Avenue,
Rockville MD, 20850

PROFESSIONAL TRAINING:

1997-1999 Post-doctoral fellow,
Advisor: James A. Johnston, Ph.D
Department of Cell Signaling,
DNAX Research Institute,
Palo Alto CA, 94304

1993-1996 Special Volunteer-Fogarty Visiting fellow,
Laboratory of Molecular Immunology
Chief: Warren J. Leonard, M.D.
National Heart, Lung, Blood Institute,
National Institutes of Health, Bethesda, MD, 20892

1992-1993 Guest Researcher,
Laboratory of Molecular Microbiology
Chief: Malcolm A. Martin, M.D.
National Institute of Allergy and Infectious Diseases,
National Institutes of Health, Bethesda, MD, 20892

EDUCATION:

- 1991-1992 Professional apprenticeship
Laboratory of AIDS-Immunopathogenesis
Chief: Professor Adriano Lazzarin
Ospedale San Raffaele, Milan, Italy
- 1991 Doctoral Degree in Biological Sciences (cum laude)
Laboratory of Macromolecular Biochemistry
Department of Biochemistry,
University of Pavia, Italy
Thesis: " Role of the oxidation state of methionine residues in
membrane proteins".

FELLOWSHIPS:

- 1993-1995 Fellowship for AIDS-related research
Istituto Superiore di Sanita',
Rome, Italy
- 1995-1996 Fogarty Visiting Fellowship,
National Institutes of Health,
Bethesda, MD 20892

PUBLICATIONS

Gianotti N, Finazzi R, Uberti Foppa C, Danise A, **Migone T**, Lazzarin A.
The thymic hormones in the treatment of immune deficiency related to HIV infection.
Pharmacol Res. 1992 Sep;26 Suppl 2:62-3.

Morsica G, Panese S, Novati R, **Migone TS**, Barin P, Mustara D, Frigato A, Lazzarin A.

Hepatitis C virus infection in blood donors with indeterminate results in second-generation recombinant immunoblot assay.

Transfusion. 1994 Jun;34(6):555-6.

Vicenzi E, Dimitrov DS, Engelman A, **Migone TS**, Purcell DF, Leonard J, Englund G, Martin MA.

An integration-defective U5 deletion mutant of human immunodeficiency virus type 1 reverts by eliminating additional long terminal repeat sequences.

J Virol. 1994 Dec;68(12):7879-90.

Lin JX, **Migone TS**, Tsang M, Friedmann M, Weatherbee JA, Zhou L, Yamauchi A, Bloom ET, Mietz J, John S, Leonard WJ

The role of shared receptor motifs and common Stat proteins in the generation of cytokine pleiotropy and redundancy by IL-2, IL-4, IL-7, IL-13, and IL-15.

Immunity. 1995 Apr;2(4):331-9.

Migone TS, Lin JX, Cereseto A, Mulloy JC, O'Shea JJ, Franchini G, Leonard WJ.
Constitutively activated Jak-STAT pathway in T cells transformed with HTLV-I.
Science. 1995 Jul 7;269(5220):79-81.

Russell SM, Tayebi N, Nakajima H, Riedy MC, Roberts JL, Aman MJ, **Migone TS**, Noguchi M, Markert ML, Buckley RH, Leonard WJ.

Mutation of Jak3 in a patient with SCID: essential role of Jak3 in lymphoid development. Science. 1995 Nov 3;270(5237):797-800.

Friedmann MC, **Migone TS**, Russell SM, Leonard WJ.

Different interleukin 2 receptor beta-chain tyrosines couple to at least two signaling pathways and synergistically mediate interleukin 2-induced proliferation.

Proc Natl Acad Sci U S A. 1996 Mar 5;93(5):2077-82.

Hilbert DM, **Migone TS**, Kopf M, Leonard WJ, Rudikoff S.

Distinct tumorigenic potential of abl and raf in B cell neoplasia: abl activates the IL-6 signaling pathway.

Immunity. 1996 Jul;5(1):81-9.

Ascherman DP, **Migone TS**, Friedmann MC, Leonard WJ.

Interleukin-2 (IL-2)-mediated induction of the IL-2 receptor alpha chain gene.

Critical role of two functionally redundant tyrosine residues in the IL-2 receptor beta chain cytoplasmic domain and suggestion that these residues mediate more than Stat5 activation.

J Biol Chem. 1997 Mar 28;272(13):8704-9.

Takemoto S, Mulloy JC, Cereseto A, **Migone TS**, Patel BK, Matsuoka M, Yamaguchi K, Takatsuki K, Kamihira S, White JD, Leonard WJ, Waldmann T, Franchini G.

Proliferation of adult T cell leukemia/lymphoma cells is associated with the constitutive activation of JAK/STAT proteins.

Proc Natl Acad Sci U S A. 1997 Dec 9;94(25):13897-902

Mulloy JC, **Migone TS**, Ross TM, Ton N, Green PL, Leonard WJ, Franchini G. Human and simian T-cell leukemia viruses type 2 (HTLV-2 and STLV-2(pan-p)) transform T cells independently of Jak/STAT activation.
J Virol. 1998 May;72(5):4408-12.

Migone TS, Cacalano NA, Taylor N, Yi T, Waldmann TA, Johnston JA. Recruitment of SH2-containing protein tyrosine phosphatase SHP-1 to the interleukin 2 receptor; loss of SHP-1 expression in human T-lymphotropic virus type I-transformed T cells.
Proc Natl Acad Sci U S A. 1998 Mar 31;95(7):3845-50.

Migone TS, Rodig S, Cacalano NA, Berg M, Schreiber RD, Leonard WJ. Functional cooperation of the interleukin-2 receptor beta chain and Jak1 in phosphatidylinositol 3-kinase recruitment and phosphorylation.
Mol Cell Biol. 1998 Nov;18(11):6416-22

Cacalano NA, **Migone TS**, Bazan F, Hanson EP, Chen M, Candotti F, O'Shea JJ, Johnston JA. Autosomal SCID caused by a point mutation in the N-terminus of Jak3: mapping of the Jak3-receptor interaction domain.
EMBO J. 1999 Mar 15;18(6):1549-58.

Cohne SJ, Sanden D, Cacalano NA, Yoshimura A, Mui A, **Migone TS**, Johnston JA. SOCS-3 is tyrosine phosphorylated in response to interleukin-2 and suppresses STAT5 phosphorylation and lymphocyte proliferation.
Mol Cell Biol. 1999 Jul;19(7):4980-8.

Aman MJ*, **Migone TS***, Sasaki A, Ascherman DP, Zhu M, Soldaini E, Imada K, Miyajima A, Yoshimura A, Leonard WJ. (* The authors contributed equally to this article)

CIS associates with the interleukin-2 receptor beta chain and inhibits interleukin-2-dependent signaling.

J Biol Chem. 1999 Oct 15;274(42):30266-72.

Wu Y, Bressette D, Carrell JA, Kaufman T, Feng P, Taylor K, Gan Y, Cho YH, Garcia AD, Gollatz E, Dimke D, LaFleur D, **Migone TS**, Nardelli B, Wei P, Ruben SM, Ullrich SJ, Olsen HS, Kanakaraj P, Moore PA, Baker KP. Tumor Necrosis Factor Receptor Superfamily Member TACI is a High Affinity Receptor for TNF Family Members APRIL and BLyS.

J Biol Chem. 2000 275(45):35478-85

Nardelli B., Belvedere O, Roschke V, Moore PA, Olsen HS, **Migone TS**,
Sosonovtseva S, Carrell JA, Feng P, Giri JG, Hilbert DM.

Synthesis and release of B-lymphocyte stimulator from myeloid cells.
Blood 2001 97 (1):198-204

Kanakaraj P, **Migone TS**, Nardelli B, Ullrich S, Li y, Olsen HS, Salcedo TW,
Kaufman T, Cochrane E, Gan Y, Hilbert DM, Giri J.
BLyS binds to B cells with high affinity and induces activation of the transcription
factors NF-kB and ELF-1.
Cytokine 2001, 13(1); 25-31

Migone TS, Humbert M, Rasclé A, Sanden D, D'Andrea A., Johnston JA
The Deubiquitinating enzyme DUB-2 prolongs cytokine-induced STAT activation
and suppresses apoptosis induced by cytokine withdrawal.
Blood 2001, 98(6), 1935-41.

Migone TS ^{*}, Zhang J ^{*}, Luo X, Zhuang Li, Chen C, Hu B, Hong JS, Perry JW,
Chen SF, Zhou JXH, Cho YH, Ullrich S, Kanakaraj P, Carrell J, Boyd E, Olsen H, Hu
G, Pukac L, Ding L, Ni J, Kim S, Gentz R, Feng P, Moore PA, Ruben SM and Wei P
(* The authors contributed equally to this article)
TL1A is a Novel Ligand for DR3 and TR6/DcR3 and Functions as a T cell
Costimulator
Immunity 2002, in press